INDEX OF AUTHORS AND SUBJECTS

Volume 131

ANDERSEN, D.E., *see* KNIGHT, R.L.

BAILLIE, S.R. & MILNE, H., Movement of Eiders *Somateria mollissima* on the east coast of Britain, 321–335.

BAINES, D., The effect of improvement of upland, marginal grasslands on the breeding success of Lapwings *Vanellus vanellus* and other waders, 497–506.

BARNARD, P. & MARKUS, M.B., Male copulation frequency and female competition for fertilizations in a promiscuous brood parasite, the Pin-tailed Whydah *Vidua macroura*, 421–425.

Behaviour
Accipiter gentilis, 205–231.
Acrocephalus palustris, 403–414.
——*scirpaceus*, 250–256.
Aegolius funereus, 41–50.
Anas penelope, 1–8.
Buteo jamaicensis, 22–26.
Bycanistes subcylindricus, 567–571.
Calcarius lapponicus, 27–32.
Carduelis flammea, 27–32.
Catharacta skua, 225–237.
Cisticola juncidis, 257–262.
Diomedea exulans, 183–195.
Falco columbarius, 426–429.
——*peregrinus*, 27–32.
Fulmarus glacialis, 225–237.
Gallinula chloropus, 429–432.
Gymnorhina tibicen, 601–608.
Larus argentatus, 225–237.
——*fuscus*, 225–237.
——*marinus*, 225–237.
Lipaugus unirufus, 303–304.
Neophron percnopterus, 9–15.
Oenanthe oenanthe, 27–32.
Parus major, 415–420, 436–442.
——*montanus*, 128–134.
Phylloscopus trochilus, 263–267.
Pica pica, 141–153.
Plectrophenax nivalis, 27–32.
Podiceps cristatus, 477–486.
Prosopeia tabuensis, 238–249.
Rissa tridactyla, 225–237.
Sarcorhamphus papa, 301–303.
Sephanoides spp., 548–566.
Somateria mollissima, 321–335.
Sula bassana, 225–237.
Tringa totanus, 214–224.
Vanellus vanellus, 377–388.
Vidua macroura, 421–425.

BERTRAM, B.C.R., *see* THOULESS, C.R.

Black Guillemot breeding biology in Shetland, 507–520.

BLAKEY, J.K., *see* NORRIS, K.J.

BOAG, D.A., *see* COURT, G.S.

BOGAN, J.A., *see* NEWTON, I.

BRADLEY, D.M., *see* COURT, G.S.

Breeding
Acrocephalus paludicola, 298–300.
Aegolius funereus, 51–56.
Aquila chrysaetos, 336–348.
——*rapax*, 33–40.
Cepphus grylle, 507.
Chrysococcyx basalis, 528.
——*lucidus*, 528.
Cisticola juncidis, 257–262.
Diomedea amsterdamensis, 171–182.
—*exulans*, 183–195.
Milvus milvus, 16–21.
Motacilla alba, 578–588.
——*grandis*, 578–588.
Parus caeruleus, 268–280.
——*major*, 112–127, 268–280, 589–600.
Prosopeia tabuensis, 238–249.
Somateria mollissima, 321–335.
Stephanoaetus coronatus, 294–298.
Tringa totanus, 214–224.
Uria aalge, 85–93.
Vanellus vanellus, 497.

BREIEHAGEN, T., Nesting biology and mating system in an alpine population of Temminck's Stint *Calidris temminckii*, 389–402.

British Ornithologists' Union
Accounts 1988, 472–475.
Annual General Meeting 1989, 628–630.
Current Notes and Notices, 165–169, 316–319, 461–466, 630–633.
Report of Council 1988, 468–471.
Union Medal: citation, 466–467.

Brood parasitism
Cettia diphone, 94–98.
Chrysococcyx basalis, 528–547.
——*lucidus*, 528–547.
Cuculus canorus, 250–256.
Vidua macroura, 421–425.

BROOKE, M. DE L., Determination of the absolute visual threshold of a nocturnal seabird, the Common Diving Petrel *Pelecanoides urinatrix*, 290–294.

BROOKE, M. DE L. & DAVIES, N.B., Provisioning of nestling Cuckoos *Cuculus canorus* by Reed Warbler *Acrocephalus scirpaceus* hosts, 250–256.

BROOKER, L.C., *see* BROOKER, M.G.

BROOKER, M.G. & BROOKER, L.C., The comparative breeding behaviour of two sympatric cuckoos, Horsfield's Bronze-Cuckoo *Chrysococcyx basalis* and the Shining Bronze-Cuckoo *C. lucidus* in Western Australia: a new model for the evolution of egg morphology and host specificity in avian brood parasites, 528–547.

Cainism in the Sulidae, 609.

Clutch interval
Calidris temminckii, 389–402.

Clutch interval (*cont.*)
Gallinula chloropus, 429–432.
Parus caeruleus, —*major*, 281–289.
Clutch overlap in Great Tits, 589–600.
Clutch size
Falco peregrinus, 355–376.
Ficedula hypoleuca, 572–577.
Parus caeruleus, 268–280.
—*major*, 268–280, 589–600.
Sulidae, 609.
COLWELL, R.K., Hummingbirds of the Juan Fernández Islands: natural history, evolution and population status, 548–566.
Contamination of Peregrine eggs, 355–376.
COURT, G.S., BRADLEY, D.M. GATES, C.C. & BOAG, D.A., Turnover and recruitment in a tundra population of Peregrine Falcons. *Falco peregrinus*, 487–496.
Courtship of the Rufous Piha, 303–304.
Crowned Eagles in rainforest, 294–298.
Cuckoldry in the Great Tit, 436–442.
Cuckoos fed by Reed Warblers, 250–256.

DAVIS, J.E., *see* NEWTON, I.
DAVIS, P.E., *see* NEWTON, I.
DAWSON, A., *see* WILLIAMS, T.D.
DE LAET, J.F. & DHONDT, A.A., Weight loss of the female during her first brood as a factor influencing second brood initiation in Great Tits *Parus major* and Blue Tits *P. caeruleus*, 281–289.
DHONDT, A.A., The effect of old age on the reproduction of Great Tits *Parus major* and Blue Tits *P. caeruleus*, 268–280.
DYRCZ, A., Polygyny in the Aquatic Warbler *Acrocephalus paludicola*, 298–300.

EBBINGE, B.S., A multifactorial explanation for variation in breeding performance of Brent Geese *Branta bernicla*, 196–204.
Ecology
Accipiter gentilis, 205–231.
Acrocephalus aequinoctialis, 99–111.
Aquila chrysaetos, 336–348.
—*rapax*, 33–40.
Buteo jamaicensis, 22–26.
Calidris temminckii, 389–402.
Chrysococcyx basalis, —*lucidus*, 528–547.
Erithacus rubecula, 432–436.
Ficedula hypoleuca, 572–577.
Falco peregrinus, 27–32.
Motacilla alba, —*grandis*, 578–588.
Parus major, 112–127.
Sagittarius serpentarius, 349–354.
Sephanoides fernandensis, —*sephanoides*, 548–566.
Tringa totanus, 214–224
Vanellus vanellus, 377–388, 497–506
EDEN, S.F., The social organization of non-breeding Magpies *Pica pica*, 141–153.
EDEN, S.F., HORN, A.G. & LEONARD, M.L., Food provisioning lowers inter-clutch interval in Moorhens *Gallinula chloropus*, 429–432.
Evolution
Egg colour discrimination in Bush Warblers, 94–98.
Egg morphology and host specificity in sympatric cuckoos, 528–547.
Island hummingbird morphology and behaviour, 548–566.
Stone throwing behaviour, 9–15.
EWINS, P.J., The breeding biology of Black Guillemots *Cepphus grylle* in Shetland, 507–520.

FANSHAWE, J.H., *see* THOULESS, C.R.
Feather eating in grebes, 477–486.
FEDUCCIA, A. & VOORHIES, M.R., Miocene hawk converges on Secretarybird, 349–354.
Feeding & Food
Accipiter gentilis, 205–231.
Acrocephalus palustris, 403–414.
—*scirpaceus*, 250–256.
Aegolius funereus, 51–56.
Anas penelope, 1–8.
Aquila chrysaetos, 336–348.
Buceros rhinoceros, 521–527.
Bycanistes subcylindricus, 567–577.
Catharacta skua, 225–237.
Chalcophaps indica, 521–527.
Charadrius semipalmatus, 69–79.
—*wilsonia*, 69–79.
Criniger phaeocephalus, 521–527.
Ducula badia, 521–527.
Falco peregrinus, 355–376.
Fulmarus glacialis, 225–237.
Gallinula chloropus, 429–432.
Gracula religiosa, 521–527.
Himantopus h. mexicanus, 69–79.
Larus argentatus, —*fuscus*, —*marinus*, 225–237.
Limnodromus griseus, 69–79.
Megalaima haemocephala, 521–527.
Oriolus chinensis, 521–527.
Parus major, 415–420.
Ptilinopus jambu, 521–527.
Pycnonotus brunneus, 521–527.
—*erythrophthalmos*, 521–527.
Rissa tridactyla, 225–237.
Stephanoaetus coronatus, 294–298.
Sula bassana, 225–237.
Treron spp., 521–527.
Tringa flavipes, 69–79.
—*melanoleuca*, 69–79.
Vanellus vanellus, 80–84, 377–388.
Feeding site selection by Wigeon, 1–8.
Flying behaviour and habitat preferences of King Vultures, 301–303.
FULLER, M.R., *see* MEESE, R.J.
FURNESS, R.W., *see* HUDSON, A.W.

GALBRAITH, H., The diet of Lapwing *Vanellus vanellus* chicks on Scottish farmland, 80–84.
GALBRAITH, H., Arrival and habitat use by Lapwings *Vanellus vanellus* in the early breeding season, 377–388.
GATES, C.C., *see* COURT, G.S.
GIBBONS, D.W., Seasonal reproductive success of the Moorhen *Gallinula chloropus*: the importance of male weight, 57–68.
GREEN, P.T. & THEOBALD, C.M., Sexing birds by discriminant analysis: further considerations, 442–447.
GREEN, R.E., Transformation of crude proportions of nests that are successful for comparison with Mayfield estimates of nest success, 305–306.

HAAS, M.B., *see* NEWTON, I.
Habitat preference of Tawny Eagles in the tropics, 33–40.
Habitat use by Lapwings, 377–388.
HALE, W.G., *see* THOMPSON, P.S.
HARRIS, M.P., Variation in the correction factor used for converting counts of individual Guillemots *Uria aalge* into breeding pairs, 85–93.
HEDENSTRÖM, A., Which wing-index should be used?, 154.
HIGUCHI, H., Responses of the Bush Warbler *Cettia diphone* to artificial eggs of *Cuculus* cuckoos in Japan, 94–98.
HIGUCHI, H. & HIRANO, T., Breeding season, courtship behaviour, and territoriality of White and Japanese Wagtails *Motacilla alba* and *M. grandis*, 578–588.
HIRANO, T., *see* HIGUCHI, H.
HOELZEL, A.R., Territorial behaviour of the Robin *Erithacus rubecula*: the importance of vegetation density, 432–436.
HOGSTAD, O., Subordination in mixed-age bird flocks—a removal study, 128–134.
HOGSTAD, O., The presence of non-territorial males in Willow Warbler *Phylloscopus trochilus* populations—a removal study, 263–267.
HORN, A.G., *see* EDEN, S.F.
HOWELLS, W.W., *see* HUSTLER, K.
HUDSON, A.V. & FURNESS, R.W., The behaviour of seabirds foraging at fishing boats around Shetland, 225–237.
Hummingbirds: comparison between endemic and continental *Sephanoides* spp., 548–566.
Hunting behaviour of Goshawk, 205–231.
HUSTLER, K. & HOWELLS, W.W., Habitat preferences, breeding success and the effect of primary productivity on Tawny Eagles *Aquila rapax* in the tropics, 33–40.

JAMES, P.C., WARKENTIN, I.G. & OLIPHANT, L.W., Turnover and dispersal in urban Merlin *Falco columbarius*, 426–429.
JÄRVINEN, A., Clutch-size variation in the Pied Flycatcher *Ficedula hypoleuca*, 572–577.
JOUVENTIN, P., MARTINEZ, J. & ROUX, J.P., Breeding biology and current status of the Amsterdam Island Albatross *Diomedea amsterdamensis*, 171–182.

KALINA, J., Nest intruders, nest defence and foraging behaviour in the Black-and-White Casqued Hornbill *Bycanistes subcylindricus*, 567–571.
KÄLLANDER, H., *see* SMITH, H.G.
KELSEY, M.G., A comparison of the song and territorial behaviour of a long-distance migrant, the Marsh Warbler *Acrocephalus palustris*, in summer and winter, 403–414.
KNIGHT, R.L., ANDERSEN, D.E., BECHARD, M.J. & MARR, N.V., Geographic variation in nest-defence behaviour of the Red-tailed Hawk *Buteo jamaicensis*, 22–26.
KORPIMÄKI, E., Mating system and mate choice of Tengmalm's Owls *Aegolius funereus*, 41–50.
KORPIMÄKI, E., Breeding performance of Tengmalm's Owl *Aegolius funereus*: effects of supplementary feeding in a peak vole year, 51–56.

LAMBERT, F.R., Pigeons as seed predators and dispersers of figs in a Malaysian lowland forest, 521–527.
Lapwing breeding success, 497–506.
LEONARD, M.L., *see* EDEN, S.F.

MACE, R., The relationship between daily routines of singing and foraging: an experiment on captive Great Tits *Parus major*, 415–420.
MARKUS, M.B., *see* BARNARD, P.
MARR, N.V., *see* KNIGHT, R.L.
MARTINEZ, J., *see* JOUVENTIN, P.
Mate choice of Tengmalm's Owls, 41–50.
Mating system
Acrocephalus paludicola, 298–300.
Aegolius funereus, 41–50.
Calidris temminckii, 389–402.
MAYHEW, P. & HOUSTON, D., Feeding site selection by Wigeon *Anas penelope* in relation to water, 1–8.
MCNEIL, R., *see* ROBERT, M.
MEESE, R.J. & FULLER, M.R., Distribution and behaviour of passerines around Peregrine *Falco peregrinus* eyries in western Greenland, 27–32.
Methods
Discriminant analysis to sex birds, 442–447.
Wing-index, 154.
MILDER, S.L. & SCHREIBER, R.W., The vocalizations of the Christmas Island Warbler *Acrocephalus aequinoctialis*, an island endemic, 99–111.
MILNE, H., *see* BAILLIE, S.R.
Movement patterns of Eider, 321–335.

NELSON, B., Cainism in the Sulidae, 609.
Nest defence by hornbills, 567–571.

Nest re-use by Fan-tailed Warblers, 257–262.
Nest sites
Bycanistes subcylindricus, 567–571.
Calidris temminckii, 389–402.
Cisticola juncidis, 257–262.
Vanellus vanellus, 377–388.
Nest success rate: crude transformed to Mayfield, 305–306.
NEWTON, I., BOGAN, J.A. & HAAS, M.B., Organochlorines and mercury in the eggs of British Peregrines *Falco peregrinus*, 355–376.
NEWTON, I., DAVIS, P.E. & DAVIS, J.E., Age of first breeding, dispersal and survival of Red Kites *Milvus milvus* in Wales, 16–21.
NICHOLLS, T.J., *see* WILLIAMS, T.D.
NILSSON, J.-Å., *see* SMITH, H.G.
NORRIS, K.J. & BLAKEY, J.K., Evidence for cuckoldry in the Great Tit *Parus major*, 436–442.

Obituaries
STANLEY CRAMP, 612–614.
DESMOND NETHERSOLE-THOMPSON, 610–611
DOMINIC LOUIS SERVENTY, 448–450.
OLIPHANT, L.W., *see* JAMES, D.C.
ORELL, M., Population fluctuations and survival of Great Tits *Parus major* dependent on food supplied by man in winter, 112–127.

PAYNE, S., *see* WATSON, A.
Peregrine population dynamics, 487–496.
Photoperiod—effect on juvenile Starlings, 135–140.
PICKERING, S.P.C., Attendance patterns and behaviour in relation to experience and pair-bond formation in the Wandering Albatross *Diomedea exulans* at South Georgia, 183–195.
PIERSMA, T. & VAN EERDEN, M.R., Feather eating in Great Crested Grebes *Podiceps cristatus*: a unique solution to the problems of debris and gastric parasites in fish-eating birds, 477–486.
Pigeons as predators and dispersers of figs, 521–527.
Polygyny in the Aquatic Warbler, 298–300.
Population dynamics
Calidris temminckii, 389–402.
Diomedea amsterdamensis, 171–182.
Falco columbarius, 426–429.
——*peregrinus*, 355–376, 487–496.
Parus major, 112–127.

RAE, R., *see* WATSON, A.
Recent Ornithological Publications, 155–164, 307–315, 451–460, 615–627.
REID, S.B., Flying behaviour and habitat preferences of the King Vulture *Sarcorhamphus papa* in the western Orinoco Basin of Venezuela, 301–303.
Reproduction in old age in Great Tits and Blue Tits, 268–280.
Reproductive biology of the Red Shining Parrot, 238–249.
Reproductive strategy of *Parus* spp., 281–289.
Reproductive success
Aquila chrysaetos, 336–348.
——*rapax*, 33–40.
Branta bernicla, 196–204.
Gallinula chloropus, 57–68.
Parus major, ——*caeruleus*, } 268–280.
Prosopeia tabuensis, 238–249.
Tringa totanus, 214–224.
RINKE, D., The reproductive biology of the Red Shining Parrot *Prosopeia tabuensis* on the island of 'Eua, Kingdom of Tonga, 238–249.
ROBERT, M. & MCNEIL, R., Comparative day and night feeding strategies of shorebird species in a tropical environment, 69–79.
ROUX, J.P., *see* JOUVENTIN, P.

SCHREIBER, R.W., *see* MILDER, S.L.
Seabirds foraging at fishing boats, 225–237.
Sexual maturation and moult in Starlings, 135–140.
SKORUPA, J., Crowned Eagles *Stephanoaetus coronatus* in rainforest: observations on breeding chronology and diet at a nest in Uganda, 294–298.
SKUTCH, A.F., Courtship of the Rufous Piha *Lipaugus unirufus*, 303–304.
SMITH, H.G., KÄLLANDER, H. & NILSSON, J.-Å., The significance of clutch overlap in Great Tits *Parus major*, 589–600.
Social organization of Magpies, 141–153, 601–608.
Song and territorial behaviour of Marsh Warblers, 403–414.
Subordination in mixed-age bird flocks, 128–134.
Supplementary feeding; effect on breeding performance of Tengmalm's Owl, 51–56.

Territoriality
Acrocephalus aequinoctialis, 99–111.
——*palustris*, 403–414.
Erithacus rubecula, 432–436.
Gymnorhina tibicen, 601–608.
Motacilla alba, ——*grandis*, } 578–588.
Phylloscopus trochilus, 263–267.
THEOBALD, C.M., *see* GREEN, P.T.
THOMPSON, P.S. & HALE, W.G., Breeding site fidelity and natal philopatry in the Redshank *Tringa totanus*, 214–224.
THOULESS, C.R., FANSHAWE, J.H. & BERTRAM, B.C.R., Egyptian Vultures *Neophron percnopterus* and Ostrich *Struthio camelus* eggs: the orgins of stone-throwing behaviour, 9–15.
Time-budget analysis
Acrocephalus palustris, 403–414.
Anas penelope, 1–8.
Charadrius semipalmatus, ——*wilsonia*, } 69–79.
Himantopus h. mexicanus, 69–79.
Parus major, 112–127, 415–420.

——*montanus*, 128–134.
Pica pica, 141–153.
Tringa melanoleuca, } 69–79.
——*flavipes*, }
Vanellus vanellus, 377–388.
Turnover and dispersal in urban Merlins, 426–429.

UEDA, K., Re-use of courtship nests for quick remating in the polygynous Fan-tailed Warbler *Cisticola juncidis*, 257–262.

VAN EERDEN, M.R., *see* PIERSMA, T.
VELTMAN, C.J., Flock, pair and group living lifestyles without cooperative breeding by Australian Magpies *Gymnorhina tibicen*, 601–608.
Vocalizations of the Christmas Island Warbler, 99–111.
VOORHIES, M.R., *see* FEDUCCIA, A.

WARKENTIN, I.G., *see* JAMES, P.C.
WATSON, A., PAYNE, S. & RAE,R., Golden Eagles *Aquila chrysaetos*: land use and food in northeast Scotland, 336–348.
Weight loss in brooding tits, 281–289.
WIDEN, P., The hunting habitats of Goshawks *Accipiter gentilis* in boreal forests of central Sweden, 205–231.
WILLIAMS T.D., DAWSON, A. & NICHOLLS, T.J., Sexual maturation and moult in juvenile Starlings *Sturnus vulgaris* in response to different daylengths, 135–140.
Wing-index: which should be used?, 154.

INDEX OF SCIENTIFIC NAMES OF BIRDS

Each name is indexed under its genus and species and, if trinomial, also under its subspecies

aalge, Uria, 85.
Acanthiza apicalis, 546.
——inornata, 528.
——pusilla, 546.
——chrysorrhoa, 528.
Accipiter cooperii, 371.
——gentilis, 33, 151, 205.
——nisus, 30, 33, 41, 51, 123, 386, 428, 435, 493.
Acrocephalus aequinoctialis, 99.
——arundinaceus, 99, 255.
——dumetorum, 412.
——paludicola, 298.
——palustris, 99, 403.
——schoenobaenus, 412.
——scirpaceus, 250, 412.
Acrocephalus spp. 300.
Actitis hypoleucos, 83, 221.
——macularia, 214, 400.
adeliae, Pygoscelis, 183.
Aegolius funereus, 41.
aequinoctialis, Acrocephalus, 99.
africanus, Hieraeetus, 295.
Alauda arvensis, 81.
alba, Calidris, 389.
——Motacilla, 578.
albicilla, Haliaeetus, 370.
albifrons, Anser, 202.
Alca torda, 85.
Alectoris rufus, 81.
alpestris, Eremophila, 489.
alpina, Calidris, 214, 504.
aluco, Strix, 293, 435.
Amazona spp., 247.
americana, Recurvirostra, 76.
americanus, Numenius, 222.
amsterdamensis, Diomedea, 171.
Anas platyrhynchos, 372.
Anser albifrons, 202.
——c. caerulescens, 333.
Anthus spinoletta, 489.
Apatosagittarius terrenus, 349.
apicalis, Acanthiza, 546.
Aplonis tabuensis, 247.
apricaria, Pluvialis, 504.
Aquila chrysaetos, 336.
——rapax, 33.
arctica, Fratercula, 21, 229.
Arenaria interpres, 69.
argentatus, Larus, 225.
aristotelis, Phalacrocorax, 229.
arquata, Numenius, 497.
arundinaceus, Acrocephalus, 99, 255.
arvensis, Alauda, 81.
ater, Molothrus, 254.
atratus, Coragyps, 301.
atricapillus, Parus, 152.
auritus, Podiceps, 485.
Aythya spp., 479.

badia, Ducula, 523.
basalis, Chrysococcyx, 528.
bassana, Sula, 193, 225.
bernicla, Branta, 196.
——, Branta bernicla, 196.
Bonasa umbellus, 30.
brachyrhynchos, Corvus, 426.
Branta b. bernicla, 196.
——leucopsis, 202.
brunneus, Pycnonotus, 525.
Bubo virginianus, 426.
bucephalus, Lanius, 97.
Buceros rhinoceros, 525.
Buteo buteo, 386.
——jamaicensis, 22.
——polyosoma, 554.
buteo, Buteo, 386.
Bycanistes subcylindricus, 567.

caerulescens, Anser c., 333.
caeruleus, Parus 268, 281.
Calcarius lapponicus, 27, 489.
Calidris alba, 389.
——alpina, 214, 504.
——maura, 214.
——minuta, 389.
——pusilla, 69, 221.
——temminckii, 214, 389.
californicus, Larus, 67.
Calyptorhynchus spp., 247.
camelus, Struthio, 9.
canorus, Cuculus, 250.
capensis, Zonotrichia, 150, 266.
caprius, Chrysococcyx, 535.
Carduelis flammea, 27.
Catharacta skua, 225, 290.
Cathartes spp., 303.
Cepphus grylle, 507.
Cettia diphone, 94.
Chalcophaps indica, 523.
chalybeata, Vidua, 424.
Charadrius semipalmatus, 69.
——wilsonia, 69.
chinensis, Oriolus, 525.
chloris, Halcyon, 247.
chloropus, Gallinula, 57, 429.
Chrysococcyx caprius, 535.
chrysaetos, Aquila, 336.
chrysorrhoa, Acanthiza, 528.
Chrysococcyx basalis, 528.
——lucidus, 528.
——l. plagosus, 529.
Cisticola juncidis, 257.
Climacteris erythrops, 535.
colchicus, Phasianus, 212, 372.
Columba livia, 293.
——palumbus, 211.
columbarius, Falco, 370, 426.
cooperii, Falco, 371.
Coragyps atratus, 301.
corax, Corvus, 398.
cornix, C. corone, 211.
coronatus, Stephanoaetus, 294.
corone, Corvus, 150, 211.
Corvus brachyrhynchos, 426.
——corax, 398.
——corone, 150.
——c. cornix, 211.
——frugilegus, 442.
——monedula, 442.
costaricensis, Zonotrichia capensis, 150.
Criniger phaeocephalus, 525.
cristatus, Podiceps, 477.
cucullatus, Ploceus, 97.
Cuculus canorus, 250.
——pallidus, 543.
Cuculus spp., 94.
cyanea, Passerina, 437.

dabbenena, Diomedea exulans, 179.
Diomedea amsterdamensis, 171.
——e. epomophora, 179.
——e. sanfordi, 179.
——exulans, 171, 183.
——e. dabbenena, 179.
diphone, Cettia, 94.
domesticus, Passer, 597.
Ducula badia, 523.
dumetorum, Acrocephalus, 412.

epomophora, Diomedea, 174.
Eremophila alpestris, 489.
Erithacus rubecula, 432.
erythrophthalmos, Pycnonotus, 525.

erythrops, Climacteris, 535.
estella, Oreotrochilus, 550.
Eudynamis scolopacea, 543.
exulans, Diomedea, 171, 183.

Falco columbarius, 370, 426.
——mexicanus, 371, 493.
——peregrinus, 355, 428, 487.
——p. tundrius, 488.
——sparverius, 370, 554.
——tinnunculus, 41, 51, 386.
fernandensis, Sephanoides, 548.
Ficedula hypoleuca, 47, 437, 572.
flammea, Carduelis, 27.
flavipes, Tringa, 69.
flavus, Lichenostomus, 535.
Fratercula arctica, 21, 229.
frugilegus, Corvus, 442.
Fulmarus glacialis, 21, 225, 370.
funereus, Aegolius, 41.
fuscus, Larus, 225.

Gallinago gallinago, 497.
gallinago, G., 497.
Gallinula chloropus, 57, 429.
Garrulus glandarius, 435.
gentilis, Accipiter, 30, 151, 205.
glacialis, Fulmarus, 21, 225, 370.
glandarius, Garrulus, 435.
Gracula religiosa, 525.
grandis, Motacilla, 578.
grisegena, Podiceps, 484.
griseus, Limnodromus, 69.
grylle, Cepphus, 507.
guttata, Poephila, 425.
Gymnorhina tibicen, 601.

haemocephala, Mesalaima, 525.
Haemotopus ostralegus, 70.
Halcyon chloris, 247.
Haliaeetus albicilla, 370.
——leucocephalus, 371.
Harpia harpyja, 296.
harpyja, Harpia, 296.
Hieraeetus africanus, 295.
Himantopus h. mexicanus, 69.
himantopus, H., 69.
Hirundo rustica, 437.
Hydrobates pelagicus, 229.
hypoleuca, Ficedula, 47, 437, 572.
hypoleucos, Actitis, 83, 221.

indica, Chalcophaps, 523.
inornata, Acanthiza, 528.
interpres, Arenaria, 69.

jamaicensis, Buteo, 22.
jambu, Ptilinopus, 523.
japonica, Zosterops, 95.
jefferyi, Pithecophaga, 295.
juncidis, Cisticola, 257.

Lagopus l. scoticus, 336.
——mutus, 336, 489.
Lalage maculosa, 247.
Lanius bucephalus, 97.
lapponicus, Calcarius, 27, 489.
Larus californicus, 67.
——ridibundus, 370.
Larus spp., 225.
leucocephalus, Haliaeetus, 371.
leucopsis, Branta, 202.
——, Motacilla alba, 578.
leucorrhoa, Oceanodroma, 290.
Lichenostomus flavus, 535.
Limnodromus griseus, 69.
Lipaugus unirufus, 303.
——vociferans, 304.
livia, Columba, 293.
lobatus, Phalaropus, 221.
lucidus, Chrysococcyx, 528.
lugens, Motacilla alba, 578.

macroura, Vidua, 421.
macularia, Actitis, 214, 400.
maculosa, Lalage, 247.
maderaspatensis, Motacilla, 578.
magnus, Pelargopappus, 353.
major, Parus, 41, 66, 112, 129, 268, 281, 415, 430, 436, 589.
Malurus splendens, 528.
marinus, Larus, 225.
maura, Calidris, 214.
Melanerpes uropygialis, 567.
melanoleuca, Tringa, 69.
melodia, Melospiza, 266.
Melospiza melodia, 266.
Mesalaima haemocephala, 525.
mexicanus, Falco, 371, 493.
——, Himantopus h., 69.
Milvus milvus, 16.
milvus, M., 16.
minor, Phoenicopterus, 14.
minuta, Calidris, 389.
modularis, Prunella, 601.
mollissima, Somateria, 321.
Molothrus ater, 254.
monachus, Neophron, 13.
monedula, Corvus, 442.
montanus, Parus, 128.
Motacilla alba, 578.
——a. leucopsis, 578.
——a. lugens, 578.
——grandis, 578.
——maderaspatensis, 578.
multicolor, Petroica, 528.
mutus, Lagopus, 336, 489.

nebularia, Tringa, 221.
Neophron monachus, 13.
——percnopterus, 9.
nisus, Accipiter, 30, 41, 51, 123, 372, 428, 435.
nivalis, Plectrophenax, 27, 489.
novaehollandiae, Scythrops, 543.
Numenius americanus, 222.
——arquata, 497.
——phaeopus, 220, 504.

occipitalis, Phylloscopus, 994
Oceanodroma leucorrhoa, 290.
ochruros, Phoenicurus, 250.
Oenanthe oenanthe, 27.
oenanthe, O., 27.
onocratolus, *Pelecanus*, 14.
Oreotrochilus estella, 550.
Oriolus chinensis, 525.
ostralegus, Haemotopus, 70.

pallidus, Cuculus, 543.
paludicola, Acrocephalus, 298.
palumbus, Columba, 211.
palustris, Acrocephalus, 109, 403.
——, Telmatodytes, 257.
papa, Sarcorhamphus, 301.
parasiticus, Stercorarius, 229.
Parus atricapillus, 152.
——caeruleus, 268, 281.
——major, 41, 66, 112, 129, 268, 281, 415, 430, 436, 589.
——montanus, 128.
Passer domesticus, 597.
Passerina cyanea, 437.
pelagicus, Hydrobates, 229.
Pelargopappus magnus, 353.
——schlosseri, 353.
Pelecanoides urinatrix, 290.
Pelecanus onocrotalus, 14.
percnopterus, Neophron, 9.
Perdix perdix, 81.
perdix, P., 81.
peregrinus, Falco, 27, 353, 428, 487.
Petroica multicolor, 528.

phaeocephalus, Criniger, 525.
phaeopus, Numenius, 220, 504.
Phalacrocorax aristotelis, 229.
Phalaropus lobatus, 221.
Phasianus colchicus, 212, 372.
Phoenicopterus minor, 14.
Phoenicurus ochruros, 250.
——phoenicurus, 97.
Phylloscopus occipitalis, 94.
——trochilus, 263.
——spp., 544.
Pica pica, 54, 97, 141, 266, 435.
pica, P., 54, 97, 141, 266, 435.
Pipreola spp., 304.
Pithecophaga jefferyi, 295.
plagosus, Chrysococcyx lucidus, 529.
platyrhynchos, Anas, 372.
Plectrophenax nivalis, 27, 489.
Ploceus cucullatus, 97.
Pluvialis apricana, 504.
——squatarola, 69.
Podiceps auritus, 485.
——cristatus, 477.
——grisegena, 484.
——taczanowskii, 485.
podiceps, Podilymbus, 485.
Podilymbus podiceps, 485.
Poephila guttata, 425.
polyosoma, Buteo, 554.
pomarinus, Stercorarius, 356.
porphyraceus, Ptilinopus, 247.
Porphyrio porphyrio, 601.
porphyrio, P., 601.
Probosciger spp., 247.
Prosopeia tabuensis, 238.
Prunella modularis, 601.
Ptilinopus jambu, 523.
——porphyraceus, 247.
Puffinus puffinus, 255, 290.
puffinus, P., 255, 290.
pusilla, Acanthiza, 546.
——, Calidris, 221.
——, Charadrius, 70.
Pycnonotus brunneus, 525.
——erythrophthalmos, 525.
Pygoscelis adeliae, 183.

rapax, Aquila, 33.
Recurvirostra americana, 76.
religiosa, Gracula, 525.
rhinoceros, Buceros, 525.
ridibundus, Larus, 370.
Rissa tridactyla, 41, 183, 225.
rolland, Rollandia, 485.
Rollandia rolland, 485.
rubecula, Erithacus, 432.
rufus, Alectoris, 81.
rustica, Hirundo, 437.

Sagittarius serpentarius, 349.
sanfordi, Diomedea epomophora, 179.
Sarcorhamphus papa, 301.
Saxicola torquata, 416.
schlosseri, Pelargopappus, 353.
schoenobaenus, Acrocephalus, 412.
scirpaceus, Acrocephalus, 250, 412.
scolopacea, Eudynamis, 543.
scoticus, Lagopus l., 336.
Scythrops novaehollandiae, 543.
semipalmatus, Charadrius, 69.
Sephanoides fernandensis, 548.
——sephaniodes, 548.
serpentarius, Sagittarius, 349
skua, Catharacta, 225, 290.
Somateria mollissima, 321.
sparverius, Falco, 370, 554.
spinoletta, Anthus 489.
splendens, Malurus, 528.
squatarola, Pluvialis, 69.
Stephanoaetus coronatus, 294.
Stercorarius parasiticus, 229.
——pomarinus, 236.
Strix aluco, 293, 435.
Struthio camelus, 9.
Sturnus vulgaris 54, 141.
subcylindricus, Bycanistes, 567.
Sula bassana, 193, 225.
Sulidae 609.

tabuensis, Aplonis, 247.
——, Prosopeia, 238.
taczanowskii, Podiceps, 485.
Telmatodytes palustris, 257.
temminckii, Calidris, 214, 386.
terrenus, Apatosagittarius, 349.
Tetrao tetrix, 211.
—— urogallus, 211.
tetrix, Tetrao, 211.
tibicen, Gymnorhina, 601.
tinnunculus, Falco, 41, 51, 386.
torda, Alca, 85.
torquata, Saxicola, 416.
totanus, Tringa, 76, 214, 497.
tridactyla, Rissa, 41, 183, 225.
Tringa flavipes, 69.
——melanoleuca, 69.
——nebularia, 221.
——totanus, 76, 214, 497.
trochilus, Phylloscopus, 263.
Troglodytes troglodytes, 250, 257.
troglodytes, T., 250, 257.
tundrius, Falco peregrinus, 488.
Turdus spp., 251.

umbellus, Bonasa, 30.
unirufus, Lipaugus, 303.
Uria aalge, 85.
urinatrix, Pelecanoides, 290.
urogallus, Tetrao, 211.
uropygialis, Melanerpes, 567.

Vanellus vanellus,76, 80, 377, 497.
vanellus, V., 76, 80, 377, 497.
Vidua chalybeata, 424.
——macroura, 421.
virginianus, Bubo, 426.
vociferans, Lipaugus, 304.
vulgaris, Sturnus, 54, 141.

wilsonia, Charadrius, 69.

Zonotrichia capensis, 266.
——c. costaricensis, 150.
Zosterops japonica, 95.